

**Table 3. Estimated distribution of adults and adolescents with a diagnosis of AIDS after a diagnosis of HIV infection, by race/ethnicity and sex, 2003—35 areas with confidential name-based HIV infection reporting**

	AIDS diagnoses				Total No.
	<12 months		≥12 months		
	No.	% <sup>a</sup>	No.	% <sup>a</sup>	
White, not Hispanic					
Male	3,647	39	5,692	61	9,399
Female	580	34	1,142	66	1,722
<b>Total</b>	<b>4,227</b>	<b>38</b>	<b>6,834</b>	<b>62</b>	<b>11,061</b>
Black, not Hispanic					
Male	5,019	42	6,916	58	11,935
Female	2,555	35	4,658	65	7,213
<b>Total</b>	<b>7,575</b>	<b>40</b>	<b>11,574</b>	<b>60</b>	<b>19,149</b>
Hispanic					
Male	2,227	41	3,233	59	5,460
Female	553	35	1,017	65	1,571
<b>Total</b>	<b>2,780</b>	<b>40</b>	<b>4,251</b>	<b>60</b>	<b>7,031</b>
Asian/Pacific Islander					
Male	112	40	171	60	284
Female	26	32	55	68	82
<b>Total</b>	<b>139</b>	<b>38</b>	<b>227</b>	<b>62</b>	<b>365</b>
American Indian/Alaska Native					
Male	54	42	76	58	130
Female	12	21	45	79	56
<b>Total</b>	<b>66</b>	<b>35</b>	<b>120</b>	<b>65</b>	<b>187</b>
<b>Total cases<sup>b</sup></b>	<b>14,842</b>	<b>39</b>	<b>23,096</b>	<b>61</b>	<b>37,938</b>

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

Data include persons in whom AIDS has developed and persons whose first diagnosis of HIV infection and the diagnosis of AIDS were made at the same time.

Since 2000, the following 35 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, Guam, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting for new diagnoses only.

<sup>a</sup> Percentages represent proportions of the total number of diagnoses of HIV/AIDS made during 2003 for the corresponding group (see row entries). Because row totals were calculated independently of the values for the subpopulations, the values in each row may not sum to the row total.

<sup>b</sup> Includes persons of unknown race or multiple races. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.